

Imaging Prostate Cancer: Clinical Utility of Prostate-Specific Membrane Antigen

David Kuppermann,¹ Jeremie Calais² and Leonard S. Marks^{1,*}

¹Department of Urology, David Geffen School of Medicine, University of California at Los Angeles, Los Angeles, California

²Ahmanson Translational Theranostics Division, Department of Molecular and Medical Pharmacology, David Geffen School of Medicine, UCLA, Los Angeles, California

Purpose: Our goal was to review the pathway and pertinent materials leading to approval of prostate-specific membrane antigen (PSMA) scanning by the U.S. Food and Drug Administration (FDA).

Materials and Methods: Beginning with the pivotal trials and working backward, we summarize the evolution of PSMA scanning, beginning with the discovery of the molecule, the mechanism of action to identify prostate cancer, the route to the present-day test and some of the major publications leading to each step of the sequence. From the thousands of PSMA articles listed on PubMed®, the present review is focused on the 4 large U.S. trials incorporating university studies of the gallium-68 compound and commercial studies of the fluorine-18 compound. The review further focuses on the role of PSMA scanning for both initial staging of prostate cancer and diagnosis of recurrent prostate cancer.

Results: PSMA is a transmembrane-bound glycoprotein which is overexpressed by 100–1,000-fold in prostate cancer cells. Preclinical PSMA studies at Cornell and Johns Hopkins in the 1990s were followed by early human studies in Germany in the early 2010s, then pivotal clinical trials at University of California, Los Angeles and University of California, San Francisco, leading to the first FDA approval in December 2020 (⁶⁸Ga-PSMA-11). In January 2021, a commercially available product (¹⁸F-DCFPyL) was approved on the basis of multisite registration trials (CONDOR and OSPREY). Sensitivity and specificity of PSMA scanning exceeds that of any other imaging method currently available for initial staging of prostate cancer and diagnosis of recurrent disease. The accuracy of PSMA scanning is attributed to the great image contrast (high signal-to-noise ratio), a property deriving from the high PSMA tracer uptake by prostate cancer cells. That property can be estimated quantitatively by a metric, the standardized uptake value. A follow-on PSMA compound, the theranostic lutetium-177, is currently pending FDA approval for treatment of metastases.

Conclusions: PSMA scanning is a disruptive technology that promises to transform the way prostate cancer is initially staged, recurrence is diagnosed and some advanced cases are treated.

Abbreviations and Acronyms

BCR = biochemical recurrence
 CT = computerized tomography
 FDA = U.S. Food and Drug Administration
 mpMRI = multiparametric magnetic resonance imaging
 MRI = magnetic resonance imaging
 NCCN® = National Comprehensive Cancer Network®
 PCa = prostate cancer
 PET = positron-emission tomography
 PPV = positive predictive value
 PSA = prostate specific antigen
 PSMA = prostate-specific membrane antigen
 RLT = radioligand therapy
 SUV = standardized uptake value
 UCLA = University of California, Los Angeles
 UCSF = University of California, San Francisco

Accepted for publication January 19, 2022.

Support: None.

Conflict of Interest: Dr. Marks is a consultant advisor to Exact Imaging and co-founder of Avenda Health; Dr. Calais reports prior consulting activities for Advanced Accelerator Applications, Blue Earth Diagnostics, Curium Pharma, GE Healthcare, IBA Radiopharma, Janssen Pharmaceuticals, Progenics Radiopharmaceuticals, Radiomedix and Telix Pharmaceuticals, outside of the submitted work. Dr. Calais is supported by the PCF (2020 Young Investigator Award 20YOUN05).

Ethics Statement: In lieu of a formal ethics committee, the principles of the Helsinki Declaration were followed.

* Correspondence: Department of Urology, David Geffen School of Medicine, University of California at Los Angeles, Wasserman Building, 300 Stein Plaza, 3rd floor, Los Angeles, California 90095 (telephone: 310-794-3070; FAX: 310-794-0987; email: lmarks@mednet.ucla.edu).

Key Words: prostate, prostatic neoplasms, glutamate carboxypeptidase, positron-emission tomography, molecular imaging

“BECAUSE of increased sensitivity and specificity, PSMA-PET/CT or PSMA-PET/MRI can serve as an...effective front-line imaging tool for initial staging and to detect biochemical recurrence of prostate cancer.”

NCCN Guidelines®, September 10, 2021¹

Prostate cancer (PCa) imaging based on prostate-specific membrane antigen (PSMA) is a disruptive innovation; its use has transformed diagnosis, staging and even treatment of PCa. PSMA is over-expressed by the great majority of PCa cells. PSMA-targeting small molecules can be bound to a radionuclide (eg gallium-68 or fluorine-18) and the conjugate can thus be visualized *in vivo* by scanning with positron-emission tomography (PET) scan. Anatomical localization of PSMA uptake is established by computerized tomography (CT) scanning overlay (PSMA PET/CT). Metastases and intra-prostatic cancer, not apparent by conventional imaging, may be seen. **Use of PSMA PET/CT has redefined the patterns of PCa spread, resulting in a profound transformation of PCa management.** PSMA PET/CT scanning was added to the National Comprehensive Cancer Network® (NCCN®) Guidelines® v1.2022 on September 10, 2021.¹ The following provides an overview of PSMA PET/CT imaging (not a systematic review), which was recently approved by the U.S. Food and Drug Administration (FDA) through use of the radiotracers ⁶⁸Ga-PSMA-11 and ¹⁸F-DCFPyL. Numerous systematic reviews (PRISMA) are available on PubMed® (75 as of January 2022). Herein we focus on the works most directly involved in the path to approval by the FDA.

BIOLOGY OF PSMA AND BASICS OF PSMA PET/CT IMAGING

PSMA (glutamate carboxypeptidase, folate hydrolyase) is a transmembrane-bound glycoprotein, native to prostate epithelial cells and overexpressed by 100–1,000-fold in PCa cells (fig. 1).² The normal function of PSMA remains unclear, but it may have a role in signaling and activating molecular pathways.³ When malignant transformation occurs, PSMA expression increases.^{4–7} The PSMA gene is located on the short arm of chromosome 11, a region not commonly deleted in PCa.⁸ The molecule has a unique 3-part structure consisting of an extracellular domain (707 amino acids), a transmembrane domain (24 amino acids) and an intracellular domain (19 amino acids; fig. 1).⁹ Since 95% of the molecule is extracellular, PSMA is readily accessible for binding of antibodies or low-molecular-weight ligands. PSMA-ligands can be labeled with radionuclides such as gallium-68 or fluorine-18 to

form a radiotracer. After injection, the PSMA-targeted radiotracer is rapidly cleared from the blood to the PSMA-binding sites, internalized and accumulated into the PSMA-expressing PCa cells.^{10,11}

Physiological PSMA expression may be seen in extraprostatic tissue, such as duodenum, small intestine, colon, kidney, salivary, lacrimal glands, nonmyelinated ganglia and the vascular supply of other cancers,¹² but not in the common sites of PCa metastasis (lymph nodes, bone).² Thus, with little background signal, allowing great contrast with PSMA-avid lesions, PET/CT visualization and localization of PCa is possible with accuracy not previously possible. A review by Maurer and associates, who were early investigators, provides extensive detail on the biology of PSMA.¹³

EVOLUTION OF PSMA-TARGETED IMAGING

Major milestones in the development of today's PSMA scan are shown in figure 2. **The proximate forerunner of today's PSMA scan was ProstaScint® (capromab pendetide/7E11-C5).**¹⁴ The ProstaScint scan (FDA approved 1996) employed a monoclonal antibody specific to the intracellular epitope of PSMA, radiolabeled with the gamma-emitter indium-111. Using gamma-camera imaging at 3 days of followup, the ProstaScint scan allowed visualization of PCa, including metastases. However, as the target was intracellular and ProstaScint could only bind to dying cells, the low signal/noise ratio reduced accuracy of ProstaScint scans. This and the poor image quality of indium-111 led to limited use and acceptance. The production of ProstaScint was discontinued in the U.S. in 2018.

In 1997, Neil Bander and others at Cornell discovered monoclonal antibodies to the extracellular epitope of PSMA (J591), conferring increased radiotracer uptake.^{15,16} At about the same time, Martin Pomper at Johns Hopkins reported that an enzyme in brain was homologous to PSMA and could be targeted with radiolabeled low-molecular weight ligands to image PCa.¹⁷ Thus was born modern interest in PSMA targeted scanning for diagnostic applications. Details of the preclinical discoveries are available elsewhere.^{18,19}

In 2012, researchers in Heidelberg, Germany reported the first patient imaged with a ⁶⁸Ga-labeled PSMA PET compound using the ligand PSMA-11.²⁰ Between 2012 and 2020 more than 1,700 studies on PSMA PET were published. On December 1st, 2020, nearly 10 years after its discovery, ⁶⁸Ga-PSMA-11 was cleared by the FDA as the first PSMA-targeted radiopharmaceutical for PET imaging of PCa.²¹ The FDA approval was originally restricted to the 2 centers where the ⁶⁸Ga-PSMA-11 radiotracer was manufactured in the pivotal trials (University of California, Los




Figure 1. Molecular structure of PSMA. The transmembrane molecule consists of 3 parts: a short intracellular domain (19 amino acids), a transmembrane domain (24 amino acids) and a long extracellular domain (707 amino acids). The extracellular domain contains an active substrate-recognition site (blue square) which is targeted by the PSMA ligands, eg ^{68}Ga -PSMA-11 and ^{18}F -DCFPyl. ^{68}Ga -PSMA-11 and ^{18}F -DCFPyl attach to the extracellular epitope of the PSMA, like the J591 antibodies. During binding, a process of internalization occurs, resulting in intracellular accumulation of the bound radioligand.¹³

Angeles [UCLA] and University of California, San Francisco [UCSF]). However, widespread availability of PSMA PET imaging is expected, following FDA

clearance of the first commercially available PSMA PET radiotracer (Pylarify®, piflufolostat-18, Progenics/Lantheus) in May 2021.²² The 4 studies of PSMA PET




Figure 2. Milestones in development of current PSMA PET/CT scan. CMS, Centers for Medicare and Medicaid Services.

Pivotal PSMA studies used in application to FDA

Indication	Studies	
BCR:		
Author	Fendler et al ²⁸	Morris et al (CONDOR) ³⁹
Yr	2019	2021
No. pts	635	208
Radiotracer	⁶⁸ Ga-PSMA-11	¹⁸ F-DCFPyL
Median age (range)	69 (44–95)	68 (43–91)
Mean ng/ml PSA (range)	2.1 (0.1–1,154)	0.8 (0.2–98.4)
% Sensitivity	90–92	Not available
% Specificity	Not available	Not available
% PPV by histopathology/correlative imaging/PSA response	84/not available/not available	93/89/100
% PPV by composite reference standard	92	Not available
% PPV by region (prostatic/pelvic/extrapelvic)	Not available	80/67/67
% Neg predictive value	Not available	Not available
Inter-reader agreement (Fleiss κ)	0.65–0.78	0.58–0.73
% Whole body detection rate	75	85–87
Pretreatment staging:		
Author	Hope et al ²⁹	Pienta et al (OSPREY) ⁴⁰
Yr	2021	2021
No. pts	277	252
Radiotracer	⁶⁸ Ga-PSMA-11	¹⁸ F-DCFPyL
Median age (range)	69 (63–73)	64 (46–84)
Mean ng/ml PSA (range)	11.1 (6.5–18)	9.7 (1.2–125.3)
% Sensitivity	40	31–42
% Specificity	95	96–99
% PPV	75	78–91
% Neg predictive value	81	81–84
Inter-reader agreement (Fleiss κ)	0.46–0.71	0.71–0.85
% Accuracy of pelvic nodal metastases detection	80	82–84

imaging, which were used in the registration trials for the 2 applications to the FDA, are shown in the table and discussed below.

Examples of PSMA PET/CT scans used for initial staging of PCa and to detect recurrence after treatment are shown in figure 3 (staging) and figure 4 (recurrence). In figure 5, the sensitivity of PSMA PET/CT for detection of metastatic PCa is contrasted with that of CT scanning alone.

GENERAL CHARACTERISTICS OF PSMA SCANS

The following are characteristics that apply universally to PSMA PET/CT scans.

Scan Performance

PET/CT scans are performed in the nuclear medicine department. Radiotracer is administered intravenously. Following tracer uptake (50–100 minutes), PET scanning is performed over the next 20–40 minutes depending on patient size and weight. Fully diagnostic CT is also performed, and the PET/CT images are fused and interpreted. The entire procedure ordinarily takes less than 2 hours including tracer uptake period.²³

Safety Profile

PSMA PET scan includes a radiation dose from the isotope of approximately 4 mSv and an average dose from the CT of 12 mSv. For adults over 50 years of age, the low-dose radiation exposure falls below concerning levels. No serious adverse effects of PSMA scanning have been reported.²⁴

PSMA PET Interpretation

Interpretation for PSMA PET imaging relies on standardized criteria (PROMISE criteria, PSMA-RADS,




Figure 3. ⁶⁸Ga-PSMA PET/CT scan (left) and corresponding CT scan (right) obtained for primary staging in a 77-year-old patient presenting with PSA of 7.1 ng/ml and prostate biopsy showing Gleason score of 4+5=9. Note clarity of cancerous lesions on PET/CT scan. PSMA expression is seen in the primary tumor (yellow arrow), lymph nodes (blue arrows) and in bone (sacrum, red arrow). NCCN Guidelines for 2021 have included PSMA PET/CT scanning for initial staging and to detect recurrence after primary treatment.¹




Figure 4. Lymph node metastases. ^{68}Ga -PSMA PET/CT scan obtained for recurrent PCa in a 68-year-old patient with PSA of 24 ng/ml 6 years following radiation therapy for Gleason score 4+4=8 PCa. Note visualization of lymph node metastases in left supraclavicular region (a), mediastinum (b), retroperitoneum (c) and pelvis (d). Radiotracers, which bind to the extracellular epitope of PSMA, allow detection of PCa at various sites throughout the body; lymph node involvement by PCa was often undetectable prior to the advent of PSMA scanning. Uptake of ^{68}Ga in the kidneys (shown here) is physiological.

E-PSMA).^{25–27} The PSMA PET reader follows a “TNM-like” structure. PSMA uptake is analyzed visually and compared to the surrounding background and organs of reference (blood, liver, salivary glands).

Standardized Uptake Value (SUV)

SUV is a metric used to estimate from the PET images the amount of radiotracer in a region. The intensity of PSMA PET signal using SUV can be graded as low (2–4.9), intermediate (5–10) and high (>10). However, SUVs cannot be used as an absolute indicator of PCa. The intensity of PSMA PET signal in a PCa lesion is related to the volume of the lesion and the level of PSMA expression which is correlated with the Gleason score.^{4–7} A high SUV (>10) observed in a location compatible with a PCa lesion on CT is highly specific for PCa. On the other hand, in lesions with low SUVs (2–4.9), there is overlap of PCa and other processes (inflammation, bone trauma, granulomatosis, ganglia, other neoplasms). As for any other PET imaging technique, the visual analysis of the signal-to-background ratio is usually more informative than the absolute intensity of the signal. The SUV represents only 1 parameter among others to make a final diagnosis and should only be used as a complement to the visual analysis.

Inter-reader agreement of interpretation of PSMA PET/CT scans among nuclear medicine physicians is substantial (kappa values 0.6–0.8).^{28,29} This is explained by the very high signal-to-noise ratio observed in PCa lesions, higher than with any other PET imaging technique.³⁰

Reproducibility of PSMA PET signal studies reported that when separate scans were performed some 2 weeks apart, the findings were nearly the same. Within-subject coefficient of variation was 12%–14% for bone and lymph node lesions,³¹ establishing that serial scanning may furnish important clinical information, ie in followup and after treatment.

Performance characteristics of PSMA PET/CT scanning are similar with both FDA-approved scans and are directly related to serum prostate specific antigen (PSA) levels (fig. 6 and text below).

DATA FROM EARLY CLINICAL TRIALS

The peer-reviewed literature on PSMA PET/CT scanning is increasing rapidly, with more than 800 publications listed on PubMed just in the year 2021. Among the first large patient cohorts gathered were those of Afshar-Oromieh³² and Eiber³³ et al in Germany. The value of PSMA PET/CT scanning has been indicated in these and subsequent large, single-arm studies.^{13,34–37} Among the universally-accepted conclusions are that PSMA PET/CT scanning 1) detects PCa lesions—including local recurrences and distant metastases (lymph nodes, bone and viscera)—with greater sensitivity than conventional imaging; 2) demonstrates great specificity for PCa; 3) retains its value during disease progression and androgen deprivation therapy, though with very low serum PSA levels detection rate decreases; 4) identifies intra-prostatic PCa; 5) provides correlation of uptake intensity with PCa disease severity and 6) paves the way for radioligand therapy (RLT) of PCa as a theranostic.




Figure 5. Increased sensitivity of PSMA PET/CT for bony lesion (arrows). CT scan (top) shows no abnormalities, but a lesion of PCa with high PSMA expression (bottom) is apparent in the right ileum. Even in retrospect, the CT scan was unrevealing. Because of the increased sensitivity and specificity, PSMA PET/CT has been added to the NCCN Guidelines for 2021.¹

PATH TO APPROVAL BY FDA

Among the clinical data most thoroughly vetted are the pivotal trials leading to recent FDA approvals (see table). The original trials, upon which the first FDA clearance was based, comprised a unique co-development effort led by imaging specialists at UCSF (Thomas Hope) and UCLA (Johannes Czernin), who were the primary investigators.^{28,29} Fendler reported the UCLA experience with biochemical recurrence (BCR); Hope reported the UCSF experience with pretreatment staging (see table); the imaging agent (⁶⁸Ga-PSMA-11) and the scanning methods were the same at both study sites. For detection of BCR (PSA >0.2 ng/ml), the scan identified the recurrence in 475/635 patients (75%); positive predictive value (PPV) by histopathological validation was 84%.²⁸ For presurgical staging, the scan identified positive pelvic lymph nodes with a sensitivity of 40% and a specificity of 95%, the low sensitivity attributable to undetected micro-metastases.²⁹

Each institution submitted similar Investigational New Drug Applications in late 2016, then matching New Drug Applications in late 2019; ⁶⁸Ga-PSMA-11 was manufactured at each place locally. On December 1, 2020 FDA clearance was received by each institution, marking the first PSMA-targeted PET/CT imaging approvals for PCa. The FDA action cleared ⁶⁸Ga-PSMA-11 for use in PET imaging for men with PCa and (1) suspected metastases who are candidates for initial therapy (2) suspected recurrence based on elevated serum PSA levels. Full details of the path to original FDA approval are available elsewhere.³⁸

FLUORINE-18 PSMA SCAN (PYLARIFY)

A second form of PSMA PET/CT scan, employing the isotope fluorine-18 with the radiotracer ¹⁸F-DCFPyL, entered U.S. clinical trials in 2016 (CONDOR) and 2018 (OSPREY) under sponsorship of Progenics, Inc. (now Lantheus following merger in 2020). Trial results were similar to those in the ⁶⁸Ga-PSMA-11 academic trials above. In the CONDOR trial of BCR (208 patients), detection rate was 59%–66% and PPV was 78%–93%.³⁹ In the OSPREY trial of metastatic/recurrent patients (93), detection rate was 96% (all biopsied) and PPV was 82%; in the OSPREY trial of men with high-risk PCa at initial staging (252), sensitivity for pelvic lymph node detection was 40% (again micro-metastases) and specificity was 98%.⁴⁰ On the basis of these data, FDA clearance was issued to Progenics for ¹⁸F-DCFPyL-PSMA in May 2021 with the same indications as ⁶⁸Ga-PSMA-11. A commercial product is now marketed as Pylarify (pifufolostat F-18).

In figure 6, performance characteristics of the gallium-68 and fluorine-18 based PSMA PET radiotracers are compared in results from separate meta-analyses.^{41,42} While a head-to-head comparison of the 2 radiotracers is not available, detection rates appear similar and are directly related to serum PSA levels. The 2 compounds ¹⁸F-DCFPyL and ⁶⁸Ga-PSMA-11 both target the extracellular domain of PSMA with great affinity, which explains their excellence as imaging agents for PCa. The main difference is the longer half-life of ¹⁸F-DCFPyL (110 minutes) vs that of ⁶⁸Ga-PSMA-11 (68 minutes), which favors supply and distribution of the commercialized product. Kit-based approaches are also available permitting on-site production of the ⁶⁸Ga-labeled compounds.

PSMA VS ¹⁸F-FLUCICLOVINE

The 2 PSMA radio-tracers for PET/CT imaging are not to be confused with the PET imaging agent ¹⁸F-fluciclovine (FACBC, Axumin, Blue Earth Diagnostics, Inc., Burlington, Massachusetts), which lacks affinity for PSMA. ¹⁸F-Fluciclovine is an amino acid analog which detects increased amino acid metabolic activity of cancer. It has been commercially available since FDA approval in 2016. A prospective




Figure 6. Performance characteristics of 2 FDA-approved versions of PSMA scanning. Dark blue bars represent ^{68}Ga -PSMA (version from UCLA and UCSF) and light blue bars represent ^{18}F -DCFPyL (version commercialized by Lantheus Corp.). Note that cancer detection rates of the 2 PSMA scans are similar and both are directly related to serum PSA levels. Data compiled from meta-analyses.^{41,42}

comparison of ^{18}F -fluciclovine (Axumin) vs ^{68}Ga -PSMA-11 PET/CT in 50 patients with early biochemical recurrent PCa after radical prostatectomy (PSA ≤ 2.0 ng/ml) found ^{68}Ga -PSMA-11 to be a superior imaging modality for overall detection rate of recurrent disease (56% vs 26%).³⁰ ^{68}Ga -PSMA-11 also had a significantly better detection rate for pelvic lymph nodes (30% vs 8%), extra-pelvic nodes (6% vs 0%), bone metastases (8% vs 0%), metastases in other organs (4% vs 0%) and extra-pelvic lesions (16% vs 0%). ^{18}F -fluciclovine was found to have a slightly better detection of lesions in the prostatic bed (18% vs 14%). The lower target-to-background ratio of fluciclovine (up to 8 times lower than for PSMA) is likely to account for differences in diagnostic performance.³⁰ Thus, PSMA appears to be the preferred PET tracer for diagnosis of recurrent PCa.

INTRAPROSTATIC LOCALIZATION OF CANCER

In addition to imaging metastases, PSMA PET/CT scanning may also help localize cancer within the

prostate. Using whole-mount pathology as a reference standard, Donato and colleagues showed an 80% overall concordance for index lesion detection between PSMA PET scans and magnetic resonance imaging (MRI).⁴³ PSMA PET/CT scans detected 13% of tumors missed by MRI; MRI detected 4% of tumors missed by PSMA PET/CT; with the combination, only 2% of index tumors were missed.⁴³ Similar results have been reported in a study employing F-18/PSMA scans.⁴⁴ Using the “gold standard” of whole-mount pathology for correlations, Sonni and colleagues recently showed that the combination of PSMA PET/CT+ multiparametric MRI (mpMRI) provided intraprostatic tumor delineation better than either modality alone.⁴⁵ Thus, pending available technology, PSMA PET/MRI or the fusion of PSMA PET/CT and mpMRI (PSMA PET+mpMRI) deserves further study for localization of primary PCa.

The potential for performing a biopsy, which targets a PSMA-avid spot, was first shown in 2015.⁴⁶ Subsequently, a fusion system was used to sample a PSMA-avid spot (PET/CT-ultrasound), thus diagnosing a




Figure 7. Example of PSMA-guided prostate biopsy in a 71-year-old patient with PSA of 8.5 ng/ml and prostate volume of 25 cc. MRI and systematic biopsy were negative (a). PSMA PET/CT scan shows 2 ^{68}Ga -PSMA-expressing lesions in prostate (b). CT overlay identifies intraprostatic lesions, which are contoured (c). Combining PET/CT with real-time ultrasound allows targeted biopsy using image-fusion device (d).⁴⁷ Biopsies of larger lesion revealed Grade Group 3 PCa.




Figure 8. RLT for advanced PCa. Survival curves (A, progression-free; B, overall) show comparison of standard treatments alone (broken lines) and standard treatments with addition of ^{177}Lu -PSMA-617 (solid lines), the radioligand targeting PSMA. Standard treatments included both androgen deprivation and taxanes in all patients. Addition of the radioligand significantly extended progression-free survival and overall survival ($p < 0.001$). Data are reconstructed from Morris et al³⁹ and Sartor et al.⁵¹

serious PCa that could not be detected otherwise.⁴⁷ An example of prostate biopsy successfully detecting an occult PCa by targeting a PSMA-avid lesion is shown in figure 7.

In a large prospective trial (PRIMARY) which compared MRI and PSMA-scanning with biopsy results, Emmett and colleagues found the 2 imaging modalities to provide additive information.³⁵ False-negative rate for clinically significant PCa was 17% with MRI and 10% with PSMA; but only 5/291 men were falsely negative by both, suggesting the

possibility of deferring biopsy in such cases. All men with SUV > 12 had clinically significant PCa.³⁵ The value of pre-biopsy imaging, as described in this important paper, deserves further study.

A PSMA-BASED THERANOSTIC

The word “theranostic” is derived from a combination of the words diagnostic and therapeutic, indicating the joint utility of an imaging agent and a therapeutic agent that both target the same receptors.

The first theranostic was radioiodine. The gamma-emission of I-131 is detectable by a scanner, and the beta-emission form provides a targeted RLT.⁴⁸ Thus theranostics allow the possibility to treat cancer lesions in a specific and tumor-selective manner.

PSMA-617 is a small molecule that, like PSMA-11, binds with high affinity to the extracellular domain of PSMA. It can be labeled with the beta-emitter lutetium-177 (^{177}Lu) for RLT. The first prospective results of ^{177}Lu -PSMA-617 therapy were reported by Hofman and associates in 2018.⁴⁹ Retrospective analysis of the German compassionate-use program determined that the RLT was effective in patients with metastatic castration-resistant PCa.⁵⁰

Sartor and colleagues confirmed safety and efficacy of ^{177}Lu -PSMA-617 therapy in a large multinational, randomized trial.⁵¹ The landmark VISION trial, which involved 831 men with metastatic castration-resistant PCa, compared standard care alone with standard care plus the RLT in men with metastases visible on ^{68}Ga -PET/CT. Treatment with ^{177}Lu -PSMA-617 was administered every 6 weeks for 4–6 cycles. When lutetium was added to standard care, significant gains were observed in progression-free survival (8.7 vs 3.4 months) and overall survival (15.3 vs 11.3 months; $p < 0.001$). Survival data from the trial are shown in figure 8. Most commonly reported side effects




Figure 9. Theranostic application of PSMA PET/CT using ^{177}Lu -PSMA in a 77-year-old patient with castration-resistant, metastatic PCa and PSA of 27 ng/ml (left). After 4 cycles of ^{177}Lu -PSMA over a 63-week period, metastases are no longer apparent (right), and PSA decreased to 0.02 ng/ml. The theranostic is a beta-emitter, which binds to the extracellular epitope of PSMA like the pure imaging agent. At the time of this writing, the PSMA theranostic is under review at FDA.

of lutetium included dry mouth, nausea and fatigue, but overall quality of life was not affected.⁵¹ With these results the sponsor Novartis filed a “new drug application” with the FDA. On September 28, 2021 the agency granted “breakthrough” status for ¹⁷⁷Lu-PSMA-617—a designation reserved for potentially transformative innovations. An example of a dramatic response to ¹⁷⁷Lu-PSMA-617 is shown in figure 9 (patient not from the trial). Approval of the therapeutic is expected in 2022.

THE FUTURE

Assuming real-world experience resembles the clinical trial outcomes, PSMA PET/CT scanning is likely to be widely adopted. A barrier to adoption is availability of the radiotracer. However, according to the manufacturer of Pylarify (Lantheus Medical Imaging, Inc., North Billerica, Massachusetts), as of January 2022 the product is available in the U.S. within a 2-hour drive of 90% of men who might require it (380 facilities, most in metropolitan areas). In the European Union and Australia, the product is available at approximately 150 sites. Another barrier might be availability of PET scanners, but currently some 2,400 PET scanners are in place in the U.S. (versus 12,000 MRI machines). The affordability barrier should come down soon, as Medicare has issued a permanent HCPCS code (Fluor-18, A9595) for the FDA-approved indications (initial staging and biochemical

recurrence), and private payors are expected to follow.

Numerous unmet PCa needs await solutions that may be provided by widespread adoption of PSMA scanning and evaluation in future clinical trials. Improved accuracy of PCa staging is one such need. Use of bone scans and abdominal CT scans would give way to the new imaging procedure, more accurate than the current staging scans. Earlier treatment for recurrent and metastatic disease would be possible, including redefinition of the oligometastatic condition. Some primary cancers, which remain undiagnosed even by MRI, would be found, especially if PET/MRI evolves; earlier curative treatments would be possible. Improved recognition of lesions suitable for partial gland ablation (or not) is another need, which is only partly resolved by current methods. The progression rate of active surveillance might be reduced, if a PSMA scan were negative up-front. For metastatic disease, the armamentarium of treatments would be expanded by therapeutics providing reduced toxicity and improved outcomes compared to what is currently available.

ACKNOWLEDGEMENTS

The current state of PSMA scanning could not have been reached without the key contributions of Thomas Hope, Ken Herrmann, Wolfgang Fendler, Matthias Eiber, Shaojun Zhu, Roger Slavik, Giuseppe Carlucci and Johannes Czernin. Mamdouh N. Aker, MSIII contributed to manuscript research. Figure 1 was created with BioRender.com.

REFERENCES

- Freedman-Cass D, Berardi R, Shead DA et al: NCCN Guidelines Version 1.2022—Prostate Cancer. Plymouth Meeting, Pennsylvania: National Comprehensive Cancer Network 2021. Available at <https://www.nccn.org/home/>.
- Silver DA, Pellicer I, Fair WR et al: Prostate-specific membrane antigen expression in normal and malignant human tissues. *Clin Cancer Res* 1997; **3**: 81.
- Cimadamore A, Cheng M, Santoni M et al: New prostate cancer targets for diagnosis, imaging, and therapy: focus on prostate-specific membrane antigen. *Front Oncol* 2018; **8**: 653.
- Kaittanis C, Andreou C, Hieronymus H et al: Correction: prostate-specific membrane antigen cleavage of vitamin B9 stimulates oncogenic signaling through metabotropic glutamate receptors. *J Exp Med* 2018; **215**: 377.
- Hupe MC, Philippi C, Roth D et al: Expression of prostate-specific membrane antigen (PSMA) on biopsies is an independent risk stratifier of prostate cancer patients at time of initial diagnosis. *Front Oncol* 2018; **8**: 623.
- Ross JS, Sheehan CE, Fisher HA et al: Correlation of primary tumor prostate-specific membrane antigen expression with disease recurrence in prostate cancer. *Clin Cancer Res* 2003; **9**: 6357.
- Perner S, Hofer MD, Kim R et al: Prostate-specific membrane antigen expression as a predictor of prostate cancer progression. *Hum Pathol* 2007; **38**: 696.
- O’Keefe DS, Su SL, Bacich DJ et al: Mapping, genomic organization and promoter analysis of the human prostate-specific membrane antigen gene. *Biochim Biophys Acta* 1998; **1443**: 113.
- Leek J, Lench N, Maraj B et al: Prostate-specific membrane antigen: evidence for the existence of a second related human gene. *Br J Cancer* 1995; **72**: 583.
- Ghosh A and Heston WDW: Tumor target prostate specific membrane antigen (PSMA) and its regulation in prostate cancer. *J Cell Biochem* 2004; **91**: 528.
- Hillier SM, Maresca KP, Femia FJ et al: Preclinical evaluation of novel glutamate-urea-lysine analogues that target prostate-specific membrane antigen as molecular imaging pharmaceuticals for prostate cancer. *Cancer Res* 2009; **69**: 6932.
- Mhaweche-Fauceglia P, Zhang S, Terracciano L et al: Prostate-specific membrane antigen (PSMA) protein expression in normal and neoplastic tissues and its sensitivity and specificity in prostate adenocarcinoma: an immunohistochemical study using multiple tumour tissue microarray technique. *Histopathology* 2007; **50**: 472.
- Maurer T, Eiber M, Schwaiger M et al: Current use of PSMA-PET in prostate cancer management. *Nat Rev Urol* 2016; **13**: 226.
- Horoszewicz JS, Kawinski E and Murphy GP: Monoclonal antibodies to a new antigenic marker in epithelial prostatic cells and serum of prostatic cancer patients. *Anticancer Res* 1987; **7**: 927.
- Smith-Jones PM, Vallabhajosula S, Goldsmith SJ et al: Vitro characterization of radiolabeled monoclonal antibodies specific for the

- extracellular domain of prostate-specific membrane antigen. *Cancer Res* 2000; **60**: 5237.
16. Tagawa ST, Vallabhajosula S, Christos PJ et al: Phase 1/2 study of fractionated dose lutetium-177-labeled anti-prostate-specific membrane antigen monoclonal antibody J591 (¹⁷⁷Lu-J591) for metastatic castration-resistant prostate cancer. *Cancer* 2019; **125**: 2561.
 17. Foss CA, Mease RC, Fan H et al: Radiolabeled small-molecule ligands for prostate-specific membrane antigen: in vivo imaging in experimental models of prostate cancer. *Clin Cancer Res* 2005; **11**: 4022.
 18. Choyke PL and Bouchelouche K: Prostate specific membrane antigen (PSMA) imaging: the past is prologue. *Translational Androl Urol* 2019; **8**: 283.
 19. Rowe SP, Gorin MA and Pomper MG: Imaging of prostate-specific membrane antigen with small-molecule PET radiotracers: from the bench to advanced clinical applications. *Annu Rev Med* 2019; **70**: 461.
 20. Eder M, Schäfer M, Bauder-Wüst U et al: ⁶⁸Ga-complex lipophilicity and the targeting property of a urea-based PSMA inhibitor for PET imaging. *Bioconjug Chem* 2012; **23**: 688.
 21. The FDA approves PSMA targeted drug for PET imaging in men with prostate cancer. *BJU Int* 2021; **127**: 267.
 22. U.S. Food and Drug Administration: FDA approves second PSMA-targeted PET imaging drug for men with prostate cancer. Available at <https://www.fda.gov/drugs/news-events-human-drugs/fda-approves-second-psma-targeted-pet-imaging-drug-men-prostate-cancer>.
 23. Fendler WP, Eiber M, Beheshti M et al: ⁶⁸Ga-PSMA PET/CT: joint EANM and SNMMI procedure guideline for prostate cancer imaging: version 1.0. *Eur J Nucl Med Mol Imaging* 2017; **44**: 1014.
 24. Jadvar H, Calais J, Fanti S et al: Appropriate use criteria for prostate-specific membrane antigen PET imaging. *J Nucl Med* 2022; **63**: 59.
 25. Eiber M, Herrmann K, Calais J et al: Prostate cancer molecular imaging standardized evaluation (PROMISE): proposed mITNM classification for the interpretation of PSMA-ligand PET/CT. *J Nucl Med* 2018; **59**: 469.
 26. Rowe SP, Pienta KJ, Pomper MG et al: PSMA-RADS version 1.0: a step towards standardizing the interpretation and reporting of PSMA-targeted PET imaging studies. *Eur Urol* 2018; **73**: 485.
 27. Ceci F, Oprea-Lager DE, Emmett L et al: E-PSMA: the EANM standardized reporting guidelines v1.0 for PSMA-PET. *Eur J Nucl Med Mol Imaging* 2021; **48**: 1626.
 28. Fendler WP, Calais J, Eiber M et al: Assessment of ⁶⁸Ga-PSMA-11 PET accuracy in localizing recurrent prostate cancer: a prospective single-arm clinical trial. *JAMA Oncol* 2019; **5**: 856.
 29. Hope TA, Eiber M, Armstrong WR et al: Diagnostic accuracy of ⁶⁸Ga-PSMA-11 PET for pelvic nodal metastasis detection prior to radical prostatectomy and pelvic lymph node dissection: a multicenter prospective phase 3 imaging trial. *JAMA Oncol* 2021; **7**: 1635.
 30. Calais J, Ceci F, Eiber M et al: ¹⁸F-fluciclovine PET-CT and ⁶⁸Ga-PSMA-11 PET-CT in patients with early biochemical recurrence after prostatectomy: a prospective, single-centre, single-arm, comparative imaging trial. *Lancet Oncol* 2019; **20**: 1286.
 31. Pollard JH, Raman C, Zakharia Y et al: Quantitative test-retest measurement of ⁶⁸Ga-PSMA-HBED-CC in tumor and normal tissue. *J Nucl Med* 2020; **61**: 1145.
 32. Afshar-Oromieh A, Avtzi E, Giesel FL et al: The diagnostic value of PET/CT imaging with the ⁶⁸Ga-labelled PSMA ligand HBED-CC in the diagnosis of recurrent prostate cancer. *Eur J Nucl Med Mol Imaging* 2015; **42**: 197.
 33. Eiber M, Maurer T, Souvatzoglou M et al: Evaluation of hybrid ⁶⁸Ga-PSMA ligand PET/CT in 248 patients with biochemical recurrence after radical prostatectomy. *J Nucl Med* 2015; **56**: 668.
 34. Hofman MS, Lawrentschuk N, Francis RJ et al: Prostate-specific membrane antigen PET-CT in patients with high-risk prostate cancer before curative-intent surgery or radiotherapy (proPSMA): a prospective, randomised, multicentre study. *Lancet* 2020; **395**: 1208.
 35. Emmett L, Buteau J, Papa N et al: The additive diagnostic value of prostate-specific membrane antigen positron emission tomography computed tomography to multiparametric magnetic resonance imaging triage in the diagnosis of prostate cancer (PRIMARY): a prospective multicentre study. *Eur Urol* 2021; **80**: 682.
 36. Roach PJ, Francis R, Emmett L et al: The impact of ⁶⁸Ga-PSMA PET/CT on management intent in prostate cancer: results of an Australian prospective multicenter study. *J Nucl Med* 2018; **59**: 82.
 37. Keidar Z, Gill R, Goshen E et al: ⁶⁸Ga-PSMA PET/CT in prostate cancer patients—patterns of disease, benign findings and pitfalls. *Cancer Imaging* 2018; **18**: 39.
 38. Carlucci G, Ippisch R, Slavik R et al: ⁶⁸Ga-PSMA-11 NDA approval: a novel and successful academic partnership. *J Nucl Med* 2021; **62**: 149.
 39. Morris MJ, Rowe SP, Gorin MA et al: Diagnostic performance of ¹⁸F-DCFPyL-PET/CT in men with biochemically recurrent prostate cancer: results from the CONDOR phase III, multicenter study. *Clin Cancer Res* 2021; **27**: 3674.
 40. Pienta KJ, Gorin MA, Rowe SP et al: A phase 2/3 prospective multicenter study of the diagnostic accuracy of prostate specific membrane antigen PET/CT with ¹⁸F-DCFPyL in prostate cancer patients (OSPREY). *J Urol* 2021; **206**: 52.
 41. Perera M, Papa N, Roberts M et al: Gallium-68 prostate-specific membrane antigen positron emission tomography in advanced prostate cancer—updated diagnostic utility, sensitivity, specificity, and distribution of prostate-specific membrane antigen-avid lesions: a systematic review and meta-analysis. *Eur Urol* 2020; **77**: 403.
 42. Treglia G, Annunziata S, Pizzuto DA et al: Detection rate of ¹⁸F-labeled PSMA PET/CT in biochemical recurrent prostate cancer: a systematic review and a meta-analysis. *Cancers* 2019; **11**: 710.
 43. Donato P, Morton A, Yaxley J et al: ⁶⁸Ga-PSMA PET/CT better characterises localised prostate cancer after MRI and transperineal prostate biopsy: is ⁶⁸Ga-PSMA PET/CT guided biopsy the future? *Eur J Nucl Med Mol Imaging* 2020; **47**: 1843.
 44. Bodar YJL, Jansen BHE, van der Voorn JP et al: Detection of prostate cancer with ¹⁸F-DCFPyL PET/CT compared to final histopathology of radical prostatectomy specimens: is PSMA-targeted biopsy feasible? The DeTeCT trial. *World J Urol* 2021; **39**: 2439.
 45. Sonni I, Felker ER, Lenis AT et al: Head-to-head comparison of ⁶⁸Ga-PSMA-11 PET/CT and mpMRI with histopathology gold-standard in the detection, intra-prostatic localization and local extension of primary prostate cancer: results from a prospective single-center imaging trial. *J Nucl Med* 2021; <https://doi.org/10.2967/jnumed.121.262398>.
 46. Zettinig O, Shah A, Hennersperger C et al: Multimodal image-guided prostate fusion biopsy based on automatic deformable registration. *Int J Comput Assist Radiol Surg* 2015; **10**: 1997.
 47. Simopoulos DN, Natarajan S, Jones TA et al: Targeted prostate biopsy using ⁶⁸Gallium PSMA-PET/CT for image guidance. *Urol Case Rep* 2017; **14**: 11.
 48. Inubushi M, Miura H, Kuji I et al: Current status of radioligand therapy and positron-emission tomography with prostate-specific membrane antigen. *Ann Nucl Med* 2020; **34**: 879.
 49. Hofman MS, Violet J, Hicks RJ et al: [¹⁷⁷Lu]-PSMA-617 radionuclide treatment in patients with metastatic castration-resistant prostate cancer (LuPSMA trial): a single-centre, single-arm, phase 2 study. *Lancet Oncol* 2018; **19**: 825.
 50. Rahbar K, Ahmadzadehfard H, Kratochwil C et al: German multicenter study investigating ¹⁷⁷Lu-PSMA-617 radioligand therapy in advanced prostate cancer patients. *J Nucl Med* 2017; **58**: 85.
 51. Sartor O, de Bono J, Chi KN et al: Lutetium-177-PSMA-617 for metastatic castration-resistant prostate cancer. *N Engl J Med* 2021; **385**: 1091.